

# Production of Complex Molecules in Astrophysical Ices

Marla H. Moore<sup>1</sup> and Reggie L. Hudson<sup>2</sup>

<sup>1</sup>Astrochemistry Laboratory, NASA/Goddard Space Flight Center,  
Greenbelt, MD 20771, USA  
email: Marla.H.Moore@nasa.gov

<sup>2</sup>Department of Chemistry, Eckerd College,  
St. Petersburg, FL 33711, USA,  
email: hudsonrl@eckerd.edu

**Abstract.** The inventory of interstellar and solar system molecules now numbers well over 100 species, including ions and molecules both charged and neutral. Gas-phase formation pathways for many of the observed organics are still uncertain, so that solid-phase syntheses are of interest. Low-temperature reactions are thought to occur within interstellar ices, on ice and grain surfaces in the interstellar medium, and on icy surfaces of solar system objects (e.g. Europa, Pluto). Ionizing radiation, such as cosmic rays, and far-UV photons are two possible initiators of such chemistry.

In the Cosmic Ice Lab at NASA's Goddard Space Flight Center, we can study both the photo- and radiation chemistries of ices at 8 - 300 K. Our most-recent work has been motivated by the detections of ethylene glycol, HOCH<sub>2</sub>CH<sub>2</sub>OH, in an interstellar source (Hollis, Lovas, Jewell, et al. 2002) and in comet Hale-Bopp (Crovisier, Bockelée-Morvan, Biver, et al. 2002). Ethylene glycol is currently the largest firmly-identified cometary molecule as well as one of the larger interstellar organics. This molecule's formation and accompanying chemistry provide challenges and tests for current astrochemical thought.

Here we discuss laboratory experiments on ethylene glycol's solid-phase formation and destruction. Using infrared spectroscopy, we have identified low-temperature radiation-chemical pathways that lead from known interstellar ices, such as either CH<sub>3</sub>OH or H<sub>2</sub>O and CO, to ethylene glycol. We also have identified a role for ethylene glycol in the formation of glycolaldehyde, HOCH<sub>2</sub>C(O)H, a simple sugar (Hollis, Lovas & Jewell 2000). Related to these results are the 6- and 7-atom molecules formed by oxidation and reduction processes in irradiated H<sub>2</sub>O + C<sub>2</sub>H<sub>2</sub> ices. Analogous experiments have been conducted on two other triply-bonded molecules in H<sub>2</sub>O-rich ice, HCN and N<sub>2</sub>. An important focus of our work is the development of reaction schemes for the formation of complex molecules within astrochemically-relevant ices, and the use of such schemes to predict new molecules awaiting detection.

**Keywords.** astrobiology - astrochemistry - ISM: abundances - ISM: molecules - methods: laboratory - molecular processes

---

## 1. Introduction

The current inventory of interstellar (IS) and solar system molecules now numbers well over a hundred species including inorganics, organics, ions, and radicals. The most complex molecule has thirteen atoms. Of the stable organics, half can be considered nitrile and acetylene derivatives, with the other half being aldehydes, alcohols, ethers, acids, ketones, amides, and related molecules. Ethylene glycol, HOCH<sub>2</sub>CH<sub>2</sub>OH, is composed of ten atoms, and is the most complex organic identified in both comet Hale-Bopp (Crovisier et al. 2002) and IS environments (Hollis et al. 2000). Table 1 lists observed interstellar and circumstellar organic molecules, with cometary species indicated by underlining.


**Table 1.** Organic molecules observed in interstellar and circumstellar regions. A few related inorganic molecules are included.

Hydrocarbons and Simple Hydrides, Oxides, and Sulfides			
$\text{CH}_4$	$\text{H}_2\text{O}$	$\text{NH}_3$	$\text{H}_2\text{S}$
$\text{C}_2\text{H}_4$	$\text{CO}$	$\text{N}_2\text{O}$	$\text{SO}_2$
$\text{C}_2\text{H}_6$	$\text{CO}_2$	$\text{N}_2$	$\text{OCS}$
Nitriles, Alkynes, and Related Molecules			
$\text{CH}_3\text{CN}$	$\text{HCN}$	$\text{HNC}$	$\text{C}_2\text{H}_2$
$\text{CH}_3(\text{CC})\text{CN}$	$\text{H}(\text{CC})\text{CN}$	$\text{CH}_3\text{NC}$	$\text{H}(\text{CC})_2\text{H}$
$\text{CH}_3(\text{CC})_2\text{CN}$	$\text{H}(\text{CC})_2\text{CN}$	$\text{HCCNC}$	$\text{H}(\text{CC})_3\text{H}$
$\text{CH}_2\text{CHCN}$	$\text{H}(\text{CC})_3\text{CN}$		$\text{CH}_3(\text{CC})\text{H}$
$\text{CH}_3\text{CH}_2\text{CN}$	$\text{H}(\text{CC})_4\text{CN}$	$\text{NH}_2\text{CN}$	$\text{CH}_3(\text{CC})_2\text{H}$
	$\text{H}(\text{CC})_5\text{CN}$		
Organics and Related Molecules			
$\text{H}_2\text{CO}$	$\text{CH}_3\text{OH}$	$\text{HCOOH}$	$\text{CH}_2\text{CO}$
$\text{HC}(\text{O})\text{CH}_3$	$\text{CH}_2\text{CH}(\text{OH})$	$\text{CH}_3\text{COOH}$	
$\text{H}_3\text{C}-\text{CH}_2-\text{C}(\text{O})\text{H}$	$\text{CH}_3\text{CH}_2\text{OH}$		$\text{HNCO}$
$\text{H}_2\text{C}=\text{CH}-\text{C}(\text{O})\text{H}$	$\text{HOCH}_2\text{CH}_2\text{OH}$	$\text{HC}(\text{O})\text{OCH}_3$	
$\text{HCCH}-\text{C}(\text{O})\text{H}$			$\text{HNCS}$
$\text{HOCH}_2\text{C}(\text{O})\text{H}$	$c\text{-C}_2\text{H}_4\text{O}$	$\text{HC}(\text{O})\text{NH}_2$	$\text{H}_2\text{CS}$
	$(\text{CH}_3)_2\text{O}$	$\text{CH}_2\text{NH}$	$\text{CH}_3\text{SH}$
$(\text{CH}_3)_2\text{CO}$		$\text{CH}_3\text{NH}_2$	

Although a variety of astrochemical problems can be approached successfully through gas-phase chemistry, efficient syntheses for many interstellar and cometary organics are still unknown, so that solid-phase reactions are of interest. Among the gas-phase interstellar observations in which solid-phase chemistry has been invoked are those involving formaldehyde,  $\text{H}_2\text{CO}$  (Federmann & Allen 1991); methanol,  $\text{CH}_3\text{OH}$  (e.g. Charnley et al. 1995, Tielens & Whittet 1997, Teixeira et al. 1998); acetic acid,  $\text{CH}_3\text{COOH}$  (Mehringer et al. 1997); ethylene oxide,  $c\text{-C}_2\text{H}_4\text{O}$  (Nummelin et al. 1998); methylenimine,  $\text{CH}_2\text{NH}$  (Dickens et al. 1997); ethyl cyanide,  $\text{CH}_3\text{CH}_2\text{CN}$  (Miao & Synder, 1997); vinyl alcohol,  $\text{CH}_2\text{CH}(\text{OH})$  (Turner & Apponi, 2001); ethylene glycol,  $\text{HOCH}_2\text{CH}_2\text{OH}$  (Hollis et al. 2002); glycolaldehyde,  $\text{HOCH}_2\text{C}(\text{O})\text{H}$  (Hollis et al. 2000); and propenal,  $\text{CH}_2\text{CHC}(\text{O})\text{H}$ , and propanal,  $\text{CH}_3\text{CH}_2\text{C}(\text{O})\text{H}$  (Hollis et al. 2004).

One way in which organics can be formed in space is through chemical reactions occurring within the icy mantles of interstellar grains. Reactions can be initiated by either particle radiation or high-energy photons, resulting in the synthesis of new molecular species which can then be ejected into the gas phase by various mechanisms (e.g. Bringa & Johnson, 2004, and references therein). Possible evidence for energetic processing within IS ices is the  $\text{OCN}^-$  ion (Hudson et al. 2001), while evidence for energetic processing in solar system ices comes from identifications of  $\text{H}_2\text{O}_2$ ,  $\text{O}_2$ , and  $\text{O}_3$  on Galilean moons. In comets, as with many interstellar regions, it is the many detections of gas-phase species that imply an active ice chemistry. In all cases, the importance of solid-phase reactions will depend on parameters such as initial ice composition, radiation flux, exposure time, temperature, and competing reactions.

Table 2 gives estimates of radiation and photon fluxes and doses for ices in both IS and solar system environments. Galactic cosmic rays are estimated to be about 87% protons,



**Figure 1.** Schematic of laboratory set-up.

12% helium nuclei, and 1% heavier nuclei, with an overall isotropic flux assumed constant for the age of the solar system,  $4.6 \times 10^9$  years. Since the IS electron cosmic-ray flux is 10-to-100 times smaller than the proton flux, for energies below 100 MeV the cosmic-ray processing of IS ices is dominated by protons. These protons have their greatest intensity in the low MeV range, and an energy spectrum decreasing with a power law index of 2.5 at higher energies. The estimated energy dose from impacting cosmic ray protons is  $\sim 0.3$  eV molecule $^{-1}$  over  $10^6$  years, the approximate lifetime of a molecular cloud. Within this same time,  $\sim 0.4$  eV molecule $^{-1}$  is deposited from UV photons that exist within dense cloud regions. Most of these IS photons are produced by hot O and B type stars, whose estimated abundance gives a calculated UV flux of  $9.6 \times 10^8$  eV cm $^{-2}$  s $^{-1}$  between 91 and 180 nm.

The final products of photon- and radiation-induced chemical reactions within ices can be, and typically are, identical, but the processes by which the reaction products are made are quite different. The energy of a 1 MeV proton is deposited in an ice along a track containing numerous excitations and ionizations, with the concomitant production of thousands of secondary electrons causing most of the observed chemical changes. In contrast, each vacuum-UV photon is absorbed in a single excitation or ionization event. For comparison, in a 1 g cm $^{-3}$  H $_2$ O-ice the range of a 1 MeV proton is  $\sim 20$   $\mu$ m, while there is a 37% drop in UV transmission in 0.15  $\mu$ m, assuming a UV cross-section of  $2 \times 10^{-18}$  cm $^2$  (Okabe 1978).

In our laboratory we synthesize complex organic molecules in cosmic-ice analogs, with reactions initiated by either high-energy protons ( $\sim 1$  MeV), to mimic cosmic rays, or vacuum-UV photons. We systematically vary reaction doses, temperatures, abundances of both minor and major (matrix) materials, and overall sample size (ice thickness). We also routinely employ isotopic labeling and isoelectronic systems to identify reaction products and test promising mechanisms. This approach allows us to understand and predict new interstellar molecules, and to identify processes and products that less comprehensive strategies would miss.

For this paper we describe the synthesis of complex organics from three triply-bonded molecules: carbon monoxide (C $\equiv$ O), acetylene (HC $\equiv$ CH), and hydrogen cyanide (HC $\equiv$ N). These unsaturated, isoelectronic, linear molecules are all highly-reactive, and all are de-

**Table 2.** Estimated UV photon and 1 MeV proton fluxes in different regions of space, and doses experienced by ices.

Region	Residence Time of Ices in Region (years)	UV <sup>a</sup>		Cosmic Ray Protons <sup>a</sup>	
		Flux 10 eV photon eV cm <sup>-2</sup> s <sup>-1</sup>	Energy Dose eV molecule <sup>-1</sup>	Flux 1 MeV proton eV cm <sup>-2</sup> s <sup>-1</sup>	Energy Dose eV molecule <sup>-1</sup>
IS Cold Dense Cloud (0.02 μm ice)	10 <sup>6</sup>	1.4×10 <sup>4</sup>	0.4	1×10 <sup>6</sup>	0.3
	10 <sup>7</sup>		4		3
Oort Cloud and KBO Region	4.6×10 <sup>9</sup>	9.6×10 <sup>8</sup>	10 <sup>8</sup> (top 0.015 μm layer)	Energy dependent flux ϕ(E)	150 (0.1 m) 55-5 (1-5 m)
Jovian Satellites e.g. Europa <sup>b</sup>	<1×10 <sup>8</sup>	4×10 <sup>13</sup>	10 <sup>13</sup> (top 0.015 μm layer)	Magnetospheric ions 7.8×10 <sup>13</sup>	150 (1 cm 10 <sup>4</sup> years)
Typical Lab Experiment	4.6×10 <sup>-4</sup>	8.6×10 <sup>14</sup>	14	7×10 <sup>10</sup>	9

<sup>a</sup>Energy dose for UV and cosmic ray protons for the IS medium is from Moore et al. (2001), for the Oort cloud, KBO and Jovian satellite regions from Moore et al. (2003) and references therein. <sup>b</sup>Possible resurfacing lifetime from Stern and McKinnon (2000).

tected in both IS and solar system environments. These three species are the starting point for making about twenty different molecules, radicals, and ions where, for CO and C<sub>2</sub>H<sub>2</sub>, the primary syntheses are based on reactions of H and OH radicals; for HCN the primary observed reaction path is oxidation of the cyano (-CN) functional group. Here we highlight radiation experiments in which we have identified the infrared (IR) signatures of complex species in H<sub>2</sub>O-rich ices containing CO, C<sub>2</sub>H<sub>2</sub>, or HCN. Since actual cosmic ices are not simple binary systems, we have also had to consider the roles played by other molecules and reactions, such as the formation of ions through acid-base chemistry with NH<sub>3</sub>. Finally, our results for these simple systems permit predictions of yet undetected, and more complex, organic molecules in comets and IS space.

## 2. Laboratory Methods

Details of our experimental set-up, ice preparation, IR spectral measurements, cryostat, and proton beam source have been published (Moore & Hudson 2000; Moore & Hudson, 1998). Figure 1 is a schematic that represents an ice sample within our sample chamber. Briefly, ices were formed by condensation of gas-phase mixtures onto a pre-cooled aluminum mirror at 10 - 20 K, where mid-IR spectra (4000 - 400 cm<sup>-1</sup>) of ices were taken by reflection from the mirror. Ices were 1 - 5 μm thick, as determined by a laser interference fringe system. Samples temperatures were controlled from 10 to 300 K with a heater on the cryostat tail section.

IR spectra were recorded before and after exposure of ice samples to a 0.8 MeV proton beam from a Van de Graaff accelerator. The use of proton irradiation to simulate cosmic-ray bombardment has been discussed earlier (e.g. Hudson & Moore, 2001; Moore et al. 1983). Since sample ice thicknesses were less than the range of the incident 0.8 MeV protons, the bombarding ions penetrated the ices and came to rest in the underlying metal substrate, not in the ices themselves. Radiation doses were determined by counting the proton fluence ( $\text{p}^+ \text{cm}^{-2}$ ) at the ice, and converting to a common scale of eV ( $18\text{-amu molecule}^{-1}$ ). Proton stopping powers and ranges were calculated with Ziegler's SRIM program (Ziegler et al. 1985; www.srim.org).

### 3. Laboratory Studies of Unsaturated Carbon-Containing Species in $\text{H}_2\text{O}$ -Rich Ices


#### 3.1. Study of CO in $\text{H}_2\text{O}$ Ice

To make a complex organic molecule, a laboratory chemist selects a simpler starting material and performs reactions such as those of addition or substitution of atoms or functional groups. In cometary and IS ices, carbon monoxide trails only  $\text{H}_2\text{O}$  in abundance, and so is a likely source of solid-phase organic chemistry. The  $\text{H}_2\text{O}$ -ice itself, subject to ionizing radiation, is a ready source of H atoms, OH radicals,  $\text{H}_2\text{O}^+$ , and  $\text{e}^-$ . These reactive species are available to participate in reactions such as  $\text{H}_2\text{O}_2$  formation, protonation by  $\text{H}_2\text{O}^+$  (or  $\text{H}_3\text{O}^+$ ), OH radical addition, and reduction by  $\text{e}^-$ . The possibility that such chemistry might take place in astronomical ices is at least 10 years old (Tielens & Allamandola 1987; Mumma et al. 1996), and probably much older.


Reaction Scheme 1 summarizes the first laboratory irradiations of  $\text{H}_2\text{O} + \text{CO}$  ices (Hudson & Moore 1999), where brackets indicate presumed but unobserved products. Radiation processing of  $\text{H}_2\text{O} + \text{CO}$  mixtures formed HCO,  $\text{H}_2\text{CO}$ , and  $\text{CH}_3\text{OH}$  through H-addition, and formic acid (HCOOH) in a branching reaction step. In addition,  $\text{CO}_2$  was efficiently made by O-atom transfer from OH radicals to CO. Figure 2 shows the IR spectrum, in the  $1900 - 1000 \text{ cm}^{-1}$  region, of an  $\text{H}_2\text{O} + \text{CO}$  (5:1) ice after irradiation to  $11 \text{ eV molecule}^{-1}$  at 16 K. Bands of  $\text{H}_2\text{CO}$ ,  $\text{CH}_3\text{OH}$ , and HCOOH in this region are identified by comparison with reference spectra of dilute mixtures of each of molecule in  $\text{H}_2\text{O}$ -ice. The HCO radical is at  $1853 \text{ cm}^{-1}$ . Similar experiments were performed at other  $\text{H}_2\text{O}/\text{CO}$  ratios (Hudson & Moore 1999) and in the presence of  $\text{NH}_3$  (Hudson et al. 2001).

One of the most important findings from these experiments was the relatively high percentage of conversion of CO into other molecules. After about  $22 \text{ eV molecule}^{-1}$ , for an ice whose initial  $\text{H}_2\text{O}/\text{CO}$  ratio was 5, the conversion of CO into  $\text{CO}_2$  was  $\sim 20\%$ , HCOOH  $\sim 40\%$ ,  $\text{H}_2\text{CO} \sim 7\%$  and  $\text{CH}_3\text{OH} \sim 9\%$ . These percentages are a function of the initial CO abundance and the radiation dose. They are calculated by knowing integrated band strengths of each molecule and assuming an ice density (Hudson & Moore 1999 for details).

A comparison of our results with hydrogenation experiments of Watanabe et al. (2003, 2004) is shown in Figure 3. In their experiments, the source of H atoms was a cold atomic hydrogen beam simulating the emplacement of H-atoms onto IS ice grain mantles in cold dark clouds. Hydrogenation of  $\text{H}_2\text{O} + \text{CO}$  (4:1) ices resulted in production of both  $\text{H}_2\text{CO}$  and  $\text{CH}_3\text{OH}$ . Watanabe et al. (2003, 2004) assumed a  $10^4 - 10^6$  year exposure to an H-atom density of  $1 \text{ cm}^{-3}$  at 10 K. Ratios of column densities of  $\text{CO}/\text{CH}_3\text{OH}$  and  $\text{H}_2\text{CO}/\text{CH}_3\text{OH}$  are plotted in Figure 3. The left-hand data point is equivalent to  $10^6$  years, and the right-hand data point to  $10^4$  years (based on Watanabe et al. 2004, Figure



REACTION SCHEME 1




**Figure 2.** New species formed in an  $\text{H}_2\text{O} + \text{CO}$  (5:1) ice irradiated to  $11 \text{ eV molecule}^{-1}$  are identified by comparison with reference spectra of dilute mixtures of organics at  $\sim 16 \text{ K}$ . Spectra are offset for clarity.

5). Our results are plotted to show data representing exposure of an  $\text{H}_2\text{O} + \text{CO}$  (5:1) ice to  $\sim 1$ , 10, and  $22 \text{ eV molecule}^{-1}$  corresponding to cosmic-ray proton exposure for  $3 \times 10^6$ ,  $3 \times 10^7$ , and  $7 \times 10^7$  years in a dense 10 K cloud. Both types of experiments appear able to convert CO to  $\text{H}_2\text{CO}$  and  $\text{CH}_3\text{OH}$  at levels comparable to those detected in cometary comae and interstellar ices. Note, however, that the rate of conversion of CO by H atoms is still controversial (Hiraoka et al. 2002, 2005).

### 3.1.1. $\text{CH}_3\text{OH}$ as a Starting Point for Ethylene Glycol Formation

Once formed,  $\text{CH}_3\text{OH}$  can undergo other reactions to make more complex molecules. Results from our experiments on  $\text{H}_2\text{O} + \text{CH}_3\text{OH}$  ice (3:1) showed that after irradiation at 16 K, CO,  $\text{CO}_2$ , HCO,  $\text{H}_2\text{CO}$ ,  $\text{HCOO}^-$ , and ethylene glycol ( $\text{HOCH}_2\text{CH}_2\text{OH}$ ) are present (Hudson & Moore, 2000a). Figure 4 shows the sample's IR spectra before and after irradiation. Absorptions at  $1088 \text{ cm}^{-1}$  (peak) and  $1046 \text{ cm}^{-1}$  (shoulder) are matched with a reference spectrum of ethylene glycol. This molecule's formation is explained by straightforward reaction chemistry in which hydroxymethyl radicals ( $\text{CH}_2\text{OH}$ ), produced by proton irradiation of  $\text{CH}_3\text{OH}$ , couple to make  $\text{HOCH}_2\text{CH}_2\text{OH}$  (schematically shown in Scheme 1).



**Figure 3.** Abundance ratios of  $\text{H}_2\text{CO}$  and  $\text{CO}$ , relative to  $\text{CH}_3\text{OH}$ , are compared for hydrogenation results of Watanabe et al. (2004) on  $\text{H}_2\text{O}+\text{CO}$  (4:1) ice at 15 K, proton irradiation results of Hudson & Moore (1999) on  $\text{H}_2\text{O} + \text{CO}$  (5:1) ice at  $\sim 16$  K, and observed ratios in interstellar ices, comets, and the Orion hot core. The graph is based on Watanabe et al. (2004, Fig. 5), and shows increasing processing from right to left. Powers of 10 indicate the ice's estimated exposure time (in years) to either the H fluence in molecular clouds (Watanabe's data) or the 1-MeV  $\text{H}^+$  cosmic-ray fluence (Hudson's data).


### 3.1.2. Relationship Between Ethylene Glycol and Glycolaldehyde

Recently we reported the solid-phase synthesis of glycolaldehyde ( $\text{HOCH}_2\text{C}(\text{O})\text{H}$ ) in irradiated ices containing ethylene glycol, both with and without  $\text{H}_2\text{O}$  (Hudson et al. 2005). This reaction is consistent with other aldehyde-alcohol interconversions which we have studied, such as for the pairs  $[\text{H}_2\text{CO}, \text{CH}_3\text{OH}]$  and  $[\text{CH}_3\text{C}(\text{O})\text{H}, \text{CH}_3\text{CH}_2\text{OH}]$  (Hudson & Moore, 2001). Figure 5 shows the changes in the IR spectrum of ethylene glycol ice during proton irradiation.


Another route we examined for glycolaldehyde formation will apply when  $\text{CH}_3\text{OH}$  exists in a CO-rich (apolar) ice. Figure 6 shows the IR spectrum of  $\text{CO} + \text{CH}_3\text{OH}$  (100:1) after radiolysis. In this ice, rupture of a CH bond gives an H atom that reacts with the CO matrix to make HCO, with a subsequent radical-radical reaction between HCO and  $\text{CH}_2\text{OH}$  giving glycolaldehyde. Two weak features in Figure 6 indicate the presence of  $\text{HOCH}_2\text{C}(\text{O})\text{H}$ , as shown by our reference spectrum. Other products include the HCO radical,  $\text{H}_2\text{CO}$ , and  $\text{CH}_3\text{C}(\text{O})\text{H}$ .

### 3.2. Study of $\text{C}_2\text{H}_2$ in $\text{H}_2\text{O}$ Ices

A second small, carbon-containing, unsaturated molecule we examined is acetylene ( $\text{C}_2\text{H}_2$ ). Acetylene forms in the photospheres of AGB stars where  $\text{C}/\text{O} \gg 1$ , locking up most of the elemental carbon. Gas-phase acetylene and its derivatives are thought to be the building blocks of PAHs and soot (e.g. Tielens & Charnley 1997). When  $\text{C}_2\text{H}_2$  is condensed into an icy grain mantle, however, H-atom addition reactions are possible, resulting in  $\text{C}_2\text{H}_6$  (Charnley et al. 1992, 1995). In fact, it was the detection of nearly equal abundances of  $\text{C}_2\text{H}_6$ ,  $\text{C}_2\text{H}_2$ , and  $\text{CH}_4$  in Comets C/1996 B2 Hyakutake and C/1996 O1 Hale-Bopp that led Mumma et al. (1996) to point out that these cometary species did not origi-




**Figure 4.** IR spectra of H<sub>2</sub>O + CH<sub>3</sub>OH ice before (upper) and after (middle) an irradiation to  $\sim 18$  eV molecule<sup>-1</sup> with 0.8 MeV protons. New products include H<sub>2</sub>CO, CH<sub>4</sub>, HCOO<sup>-</sup>, and ethylene glycol. At the bottom is a spectrum of unirradiated H<sub>2</sub>O + ethylene glycol for reference. Not shown are several other confirming bands for ethylene glycol between 500 and 1000 cm<sup>-1</sup>.




**Figure 5.** IR spectra of ethylene glycol before and after proton irradiation to a dose of  $\sim 3$ , 10, and 30 eV molecule<sup>-1</sup> compared to a spectrum of glycolaldehyde (GA) at 20 K.

nate in a thermochemically-equilibrated region of the solar nebula. Instead, the observed abundances were consistent with expected production of C<sub>2</sub>H<sub>6</sub> in interstellar icy grain mantles. Similar abundances for these organic volatiles were also found in Comet Lee, 153P/Ikeya-Zhang, and 9P/Tempel-1 (post-impact). See Mumma et al. (2005) and references therein.




**Figure 6.** IR spectra of CO + CH<sub>3</sub>OH (100:1) before (lower) and after (upper) irradiation with 0.8 MeV protons to a dose of  $\sim 1$  eV molecule<sup>-1</sup>. A reference spectrum of CO + glycolaldehyde (GA) (600:1) is compared with weak features on the wings of the large 1730 cm<sup>-1</sup> band.




#### REACTION SCHEME 2

We have shown experimentally that radiation processing of H<sub>2</sub>O + C<sub>2</sub>H<sub>2</sub> ices forms C<sub>2</sub>H<sub>4</sub> and C<sub>2</sub>H<sub>6</sub> through H addition (Moore & Hudson 1998) in the condensed phase. CH<sub>3</sub>C(O)H, C<sub>2</sub>H<sub>6</sub>, CH<sub>2</sub>CH(OH), and C<sub>2</sub>H<sub>5</sub>OH form through the multi-step reactions seen in Reaction Scheme 2. In addition, both CO<sub>2</sub> and CO are detected along with H<sub>2</sub>CO. Figure 7 shows IR spectra of H<sub>2</sub>O + C<sub>2</sub>H<sub>2</sub> (4:1) at 20 K before and after a dose of 17 eV molecule<sup>-1</sup>. Additional C<sub>2</sub>H<sub>6</sub> absorptions at 2880, 2939, and 2975 cm<sup>-1</sup> were stronger than the weak 1464 cm<sup>-1</sup> band shown in this region, and aided in the spectral assignments. The identifications of the isomers vinyl alcohol and acetaldehyde, CH<sub>2</sub>CH(OH) and CH<sub>3</sub>C(O)H respectively, are especially interesting (Hudson & Moore, 2003) as both are known interstellar molecules. Our experiments support the idea that a similar chemistry produces these molecules in interstellar space.

The abundance ratios of CH<sub>4</sub>, C<sub>2</sub>H<sub>6</sub>, and C<sub>2</sub>H<sub>2</sub> in these experiments are sufficient to explain the cometary observations. Variables that did change the final abundances include the radiation dose, the dilution of C<sub>2</sub>H<sub>2</sub> in the starting ice, and the initial presence of CH<sub>4</sub> in the ice. Since we did not observe C<sub>2</sub>H<sub>2</sub> formation in irradiated H<sub>2</sub>O + CH<sub>4</sub> mixtures, but did find CH<sub>4</sub> formation in H<sub>2</sub>O + C<sub>2</sub>H<sub>2</sub> ices, we conclude that relevant



**Figure 7.** IR spectra of  $\text{H}_2\text{O} + \text{C}_2\text{H}_2$  before and after irradiation to a dose of  $17 \text{ eV molecule}^{-1}$ . New products are identified by comparison with appropriate reference spectra (see Moore & Hudson 1998).



**REACTION SCHEME 3**


hydrocarbon chemistry for these comets is founded on H addition to  $\text{C}_2\text{H}_2$ , as opposed to  $\text{CH}_4$  reactions. This suggests that  $\text{C}_2\text{H}_2$  is part of the natal cometary ice composition.

Extrapolating these results to a more complex molecule, propynal ( $\text{HC}\equiv\text{CC}(\text{O})\text{H}$ ), similar H-addition reactions are expected to form propenal and propanal. All three of these aldehydes have been observed in Sagittarius B2(N) (Hollis et al. 2004). The H-atom addition sequence connecting these molecules is expected to be that shown in Reaction Scheme 3. Although only the first three organics in Reaction Scheme 3 have been identified as interstellar, the conversion of aldehydes to alcohols is consistently found in our experiments. Therefore, we safely predict the conversion of propenal into *n*-propanol, and encourage observational searches for the latter.


### 3.3. Study of HCN in $\text{H}_2\text{O}$ Ices

A third unsaturated, carbon-containing molecule we examined is HCN. This molecule is seen in the gas phase in the interstellar medium (e.g. Snyder & Buhl, 1971; Boonman et al. 2001) and in cometary comae (e.g. Ip et al. 1990; Bockelée-Morvan et al. 1994; Magee-Sauer et al. 1999). It is also thought that HCN is a native nuclear ice component in comets. The idea that HCN might participate in the chemistry of interstellar ice mantles was the motivation for our experiments on  $\text{H}_2\text{O} + \text{HCN}$  ices (Gerakines et al. 2004).

Solid-phase HCN reactions analogous to those with CO and  $\text{C}_2\text{H}_2$  will result in methylenimine ( $\text{CH}_2\text{NH}$ ) and methylamine ( $\text{CH}_3\text{NH}_2$ ), both being IS gas-phase molecules. However, our experiments showed that oxidation products of HCN dominated the IR spectrum of  $\text{H}_2\text{O} + \text{HCN}$  ices after radiolysis. Figure 8 shows the  $2400 - 2000 \text{ cm}^{-1}$  region



**Figure 8.** IR spectra of  $\text{H}_2\text{O} + \text{HCN}$  before and after irradiation to a dose of  $\sim 1$ , 6, 12, and 18  $\text{eV molecule}^{-1}$ . New products include HNCO,  $\text{OCN}^-$ , and  $\text{CO}_2$ .



**REACTION SCHEME 4**

for  $\text{H}_2\text{O} + \text{HCN}$  (5:1) at 18 K before and after several radiation doses (Gerakines et al. 2004). Identified products are the cyanate anion ( $\text{OCN}^-$ ), isocyanic acid (HNCO), and  $\text{CO}_2$ . Features in the 1800 - 1000  $\text{cm}^{-1}$  region (not shown) were identified with the  $\text{NH}_4^+$  cation at 1470  $\text{cm}^{-1}$  and formamide ( $\text{HCONH}_2$ ) at both 1686 and 1386  $\text{cm}^{-1}$ . Steps leading to these products are shown in Reaction Scheme 4.

Although IS ices are dominated by  $\text{H}_2\text{O}$ , and neither HCN nor any organic nitrile has been identified in them, these ices often show an intense IR feature due to  $\text{OCN}^-$  (Hudson et al. 2001). This observation of  $\text{OCN}^-$ , as opposed to HCN and nitriles, can be understood from our experiments, since our work demonstrates the ease with which  $\text{OCN}^-$  forms in  $\text{H}_2\text{O} + \text{HCN}$  mixtures. Moreover, HCN is not the only possible source of cyanate, as subsequent irradiations of many nitriles in  $\text{H}_2\text{O}$ -ice all showed  $\text{OCN}^-$  formation (Hudson & Moore 2004). This suggests that the cyanate ion may be a marker for energetic processing. Like  $\text{CO}_2$ , the cyanate ion is a stable, linear, 16-electron species.

It has been seen in warming experiments to  $\sim 200$  K (Moore & Hudson 2003), and is predicted to be on the surfaces of outer solar system bodies.

HNCO, indicated in Figure 8, is detected only as a gas-phase species in both interstellar sources and comets (e.g. Gibb et al. 2000; Schutte & Khanna 2003 and references therein). Its absence in IS ices suggests that it reacts to form  $\text{OCN}^-$  and a counter-ion. Candidate bases include  $\text{H}_2\text{O}$ , formamide, and methylamine. More likely, HNCO combines with  $\text{NH}_3$  to make  $\text{OCN}^-$  and  $\text{NH}_4^+$ . In fact, features in several interstellar sources are attributed to  $\text{NH}_4^+$  (Schutte & Khanna, 2003).

#### 4. Conclusions

Our experiments demonstrate that several types of chemical reactions are involved in making complex molecules. In irradiated  $\text{H}_2\text{O}$ -rich ices containing CO or  $\text{C}_2\text{H}_2$ , H- and OH- additions produce many stable organics, some more oxidized than the reactants, and some more reduced. Radicals formed during irradiation can dimerize, thereby lengthening carbon chains. Examples include the dimerization of  $\text{CH}_3$  to form  $\text{C}_2\text{H}_6$  (Moore & Hudson 1998) and, more recently, the dimerization of  $\text{CH}_2\text{OH}$  to make  $\text{HOCH}_2\text{CH}_2\text{OH}$  (Hudson et al. 2005). Current experiments involve studies of  $2 \text{CH}_2\text{CN} \rightarrow (\text{CH}_2\text{CN})_2$  in processed  $\text{H}_2\text{O} + \text{CH}_3\text{CN}$ . This is work in progress and relies on HPLC analysis of irradiated ice residues. It shows that identification of complex molecules are often at the limit of IR capabilities and require more-sensitive analytical techniques.

We have also found isomerization reactions to help explain the presence of isomeric pairs such as vinyl alcohol and acetaldehyde. Our work shows that vinyl alcohol is made in irradiated  $\text{H}_2\text{O} + \text{C}_2\text{H}_2$ , and then isomerizes to acetaldehyde. Among the nitriles, isomerizations include  $\text{HCN} \rightarrow \text{HNC}$  and  $\text{CH}_3\text{CN} \rightarrow \text{CH}_3\text{NC}$  (Hudson & Moore 2004), and formation of ketenimines such as  $\text{CH}_3\text{CN} \rightarrow \text{H}_2\text{C}=\text{C}=\text{NH}$ . Extensions give a prediction of interstellar carbodiimide from cyanamide ( $\text{NH}_2\text{C}\equiv\text{N} \rightarrow \text{HN}=\text{C}=\text{NH}$ ) and allene from methylacetylene ( $\text{CH}_3\text{C}\equiv\text{CH} \rightarrow \text{H}_2\text{C}=\text{C}=\text{CH}_2$ ).

Interconversions of aldehydes and alcohols are examples of both H-atom addition and the reverse reaction (hydrogen elimination). Our work has included the following:  $[\text{H}_2\text{CO}, \text{CH}_3\text{OH}]$ ,  $[\text{CH}_3\text{C}(\text{O})\text{H}, \text{CH}_3\text{CH}_2\text{OH}]$ , and  $[\text{HOCH}_2\text{C}(\text{O})\text{H}, \text{HOCH}_2\text{CH}_2\text{OH}]$ . This chemistry is expected to apply to more-complex pairs such as  $[\text{HC}\equiv\text{C}-\text{C}(\text{O})\text{H}, \text{HC}\equiv\text{C}-\text{CH}_2\text{OH}]$ ,  $[\text{H}_2\text{C}=\text{CH}-\text{C}(\text{O})\text{H}, \text{H}_2\text{C}=\text{CH}-\text{CH}_2\text{OH}]$ , and  $[\text{H}_3\text{C}-\text{CH}_2-\text{C}(\text{O})\text{H}, \text{H}_3\text{C}-\text{CH}_2-\text{CH}_2\text{OH}]$ . Therefore, our experiments suggest that IS aldehyde-alcohol pairs are likely if either member participates in condensed-phase radiation chemistry.

In all cases, formation paths compete with dissociation and elimination processes that lead to less-complex molecules. Acid-base reactions can also play a role, particularly regarding the formation of ions. Altogether, the reactions examined lead to about 50% of the known stable IS organics containing oxygen, and most of those found in comets. These observations show that discovering the identities of and reactions paths for complex interstellar molecules will require a firm understanding of simpler systems.

#### Acknowledgements

NASA funding through the Planetary Atmospheres and SARA programs, and the Goddard Center for Astrobiology, is acknowledged. R.L.H. acknowledges the support of NASA for Grant NAG-5-1843. Claude Smith and Steve Brown are thanked for their work with the Van de Graaff accelerator at the Goddard Space Flight Center.

#### References

- Allamandola, L. J., Sandford, S. A., & Valero, G. J. 1988, *Icarus* 76, 225
- Bockelée-Morvan, D., Padman, R., Davies, J. K., & Crovisier, J. 1994, *Planet. Space Sci.*, 42, 655
- Boonman, A. M. S., Stark, R., van der Tak, F. F. S., van Dishoeck, E. F., van der Wal, P. B. Schäfer, F., de Lange, G., & Laauwen, W. M. 2001, *ApJ*, 553, L63
- Bringa, E. M. & Johnson, R. E. 2004, *ApJ* 603, 159
- Charnley, S. B., Kress, M. E., Tielens, A. G. G. M., & Millar, T. J. 1995, *ApJ*, 448, 232
- Charnley, S. B., Tielens, A. G. G. M., & Millar, T. J. 1992, *ApJ*, 399, L71
- Crovisier, J., Bockelée-Morvan, D., Biver, N., Colom, P., Despois, D., & Lis, D. C. 2004, *A&A*, 418, L35
- Dickens, J. E., Irvine, W. M., & DeVries, C. H. 1997, *ApJ*, 312, 307
- Federman, S. R. & Allen, M. 1991, *ApJ*, 375, 157
- Gerakines, P. A., Moore, M. H. & Hudson, R. L. 2004, *Icarus*, 170, 204
- Gibb, E. L., Whittet, D. C. B., & Chair, J. E. 2001, *ApJ*, 558, 702
- Hiraoka, K. & 9 coauthors 2005, *ApJ*, 620, 542
- Hiraoka, K., Sato, T., Sato, S., Sogoshi, N., Yokoyama, T., Takashima, H., & Kitagawa, S. 2002, *ApJ*, 577, 262
- Hollis, J. M., Lovas, F. J., & Jewell, P. R. 2000, *ApJ*, 540, L107
- Hollis, J. M., Lovas, F. J., Jewell, P. R., & Coudert, L. H. 2002, *ApJ*, 571, L59
- Hollis, J. M., Jewell, P. R., Lovas, F. J., Remijan, A., & Møllendal, H. 2004, *ApJ*, 610, L21
- Hudson, R. L., Moore, M. H., & Cook, A. M. 2005, *Adv. Space Res.*, (in press)
- Hudson, R. L. & Moore, M. H. 2004, *Icarus*, 172, 466
- Hudson, R. L. & Moore, M. H. 2003, *ApJ*, 586, L107
- Hudson, R. L., Moore, M. H., & Gerakines, P. A. 2001, *ApJ*, 550, 1140
- Hudson, R. L. & Moore, M. H. 2001, *JGR-Planets*, 106, 33275
- Hudson, R. L. & Moore, M. H. 2000a, *Icarus*, 145, 661
- Hudson, R. L. & Moore, M. H. 2000b, *A&A*, 357, 787
- Hudson, R. L. & Moore, M. H. 1999, *Icarus*, 140, 451
- Ip, W. -H., et al. 1990, *Ann. Geophys.*, 8, 319
- Magee-Sauer, K., Mumma, M. J., DiSanti, M. A., Dello-Russo, N., & Rettig, T. W. 1999, *Icarus*, 142, 498
- Mehring, D. M., Snyder, L. E., & Miao, Y. 1997, *ApJ*, 480, L71
- Miao, Y. & Snyder, L. E. 1997, *ApJ*, 480, L67
- Moore, M. H. & Hudson, R. L. 2003b, *Icarus*, 161, 486
- Moore, M. H., Hudson, R. L., & Ferrante, R. F. 2003a, *Earth, Moon & Planets*, 92, 291
- Moore, M. H. & Hudson, R. L. 2000, *Icarus*, 145, 282
- Moore, M. H., Hudson, R. L., & Gerakines, P. A. 2001, *Spectrochimica Acta*, 57, 843
- Moore, M. H. & Hudson, R. L. 1998, *Icarus*, 135, 518
- Mumma, M. J. & 12 co-authors 2005, *Science*, (submitted)
- Mumma, M. J., DiSanti, M. A., Dello-Russo, N., Fomenkova, M., Magee-Sauer, K., Kaminski, C. D., & Xie, D. X. 1996, *Science*, 272, 1310
- Nummelin, A., Dickens, J. E., Bergman, P. Hjalmarsen, Å., Irvine, W. M., Ikeda, M., & Ohishi, M. 1998, *A&A*, 337, 275
- Okabe, H. 1978, *Photochemistry of Small Molecules*, Wiley, New York
- Schutte, W. A. & Khanna, R. K. 2003, *A&A*, 398, 1049
- Snyder, L. E. & Buhl, D. 1971, *ApJ*, 163, L47
- Teixeira, T. C., Emerson, J. P., & Palumbo, M. E. 1998, *A&A*, 330, 711
- Tielens, A. G. G. M. & Whittet, D. C. B. 1997, in *Molecules in Astrophysics: Probes and Processes*, ed., E. F. van Dishoeck, Kluwer Academic Pub., 45
- Tielens, A. G. G. M. & Charnley, S. G. 1997, in *Origins of Life and Evolution of the Biosphere*, Kluwer Academic Pub., 23
- Tielens, A. G. G. M. & Allamandola, L. J. 1987, in *Interstellar Processes*, eds. D. J. Hollenbach & H. A. Thronson, Reidel Pub., Dordrecht, 397.
- Turner, B. E. & Apponi, A. J. 2001, *ApJ*, 561, L207
- Watanabe, N., Nagaoka, A., Shiraki, R., & Kouchi A. 2004, *ApJ*, 616, L638

Watanabe, N., Shiraki, T., & Kouchi, A. 2003, *ApJ*, 588, L121

Ziegler, J. P., Biersack, J. P., & Littmark, U. 1985, *The Stopping and Range of Ions in Solids*, Pergamon Press, New York, also see: <http://www.srim.org/>

## **Discussion**